

Yaroslavl State University
Preprint YARU-HE-01/05

hep-ph/0108046

Photon-pair conversion into neutrinos in a strong magnetic field

A.V. Kuznetsov and N.V. Mikheev

*Division of Theoretical Physics, Department of Physics,
Yaroslavl State University, Sovietskaya 14,
150000 Yaroslavl, Russian Federation*

Abstract

A general analysis of the three-vertex loop amplitude in a strong magnetic field, based on the asymptotic form of the electron propagator in the field, is performed. In order to investigate the photon-neutrino process $\gamma\gamma \rightarrow \nu\bar{\nu}$, the vertex combinations of the scalar-vector-vector (SVV), pseudoscalar-vector-vector (PVV), 3-vector (VVV), and axial-vector-vector-vector (AVV) types are considered. It is shown that only the SVV amplitude grows linearly with the magnetic field strength, while in the other amplitudes, PVV , VVV , and AVV , the linearly growing terms are cancelled. The process $\gamma\gamma \rightarrow \nu\bar{\nu}$ is investigated in the left-right-symmetric extension of the standard model of electroweak interaction, where the effective scalar $\nu\bar{\nu}ee$ coupling could exist. Possible astrophysical manifestations of the considered process are discussed.

Submitted to Modern Physics Letters A

Strong magnetic fields which could be generated in the astrophysical cataclysms like a supernova explosion or a coalescence of neutron stars, make an active influence on quantum processes, thus allowing or enhancing the transitions which are forbidden or strongly suppressed in vacuum. However, the magnetic field influences significantly the quantum processes only in the case when it is strong enough. There exists a natural scale for the field strength which is the so-called critical Schwinger value $B_e = m_e^2/e \simeq 4.41 \cdot 10^{13}$ G (we use natural units in which $c = \hbar = 1$, hereafter e is the elementary charge). Among others, the set of quantum processes is very interesting where only electrically neutral particles, such as neutrinos and photons, present in the initial and final states. The external field influence on these loop processes is provided by the sensitivity of the charged virtual fermion to the field. The electron plays the main role here as the particle with the maximal specific charge, e/m_e .

In this letter we consider a three-vertex loop process in a strong magnetic field, which is described by the Feynman diagram shown in Fig.1. Here the double lines correspond to the electron propagators constructed on the base of the exact solutions of the Dirac equation in a magnetic field.

The intensively discussed process of this type is the conversion of the photon pair into the neutrino - antineutrino pair, $\gamma\gamma \rightarrow \nu\bar{\nu}$, which is strongly suppressed in vacuum. For this process, two vertices are vectors, e.g. $\Gamma_1 = \Gamma_2 = V$, and the third one can be vector and axial-vector, $\Gamma_3 = V, A$ (in the standard model), or it could be scalar and pseudoscalar, $\Gamma_3 = S, P$ (beyond the standard model). It is well-known (the so-called Gell-Mann theorem [1]), that for massless neutrinos, for both photons on-shell and in the local limit of the standard-model weak interaction, the process $\gamma\gamma \rightarrow \nu\bar{\nu}$ is forbidden. With any deviation from the Gell-Mann theorem conditions, the process becomes allowed, e.g. for finite neutrino mass [2, 3] or with the non-locality of the weak interaction taken into account [4].

This process becomes also possible, in the frame of the standard model, in the presence of external magnetic field. In the limit of relatively weak magnetic field, $B \ll B_e$, the process was considered in Refs. [5]. Our aim is to investigate the process $\gamma\gamma \rightarrow \nu\bar{\nu}$ in the strong field limit, both in the standard model and beyond. So, it is worthwhile to set Γ_1 and Γ_2 as vector vertices contracted with photons, keeping Γ_3 to be arbitrary.

The electron propagator in a magnetic field can be presented in the form

$$S(x, y) = e^{i\Phi(x, y)} \hat{S}(x - y), \quad (1)$$

$$\Phi(x, y) = -e \int_x^y d\xi_\mu \left[A_\mu(\xi) + \frac{1}{2} F_{\mu\nu}(\xi - y)_\nu \right], \quad (2)$$

where A_μ is a 4-potential of the external uniform magnetic field. The translational invariant part $\hat{S}(x - y)$ of the propagator has several representations. We take its asymptotic form in the strong field limit, when the magnetic field strength is the maximal physical parameter:

$$\hat{S}(X) \simeq \frac{ieB}{2\pi} \exp\left(-\frac{eBX_\perp^2}{4}\right) \int \frac{d^2p}{(2\pi)^2} \frac{(p\gamma)_\parallel + m}{p_\parallel^2 - m^2} \Pi_- e^{-i(pX)_\parallel}, \quad (3)$$

$$d^2p = dp_0 dp_3, \quad \Pi_- = \frac{1}{2}(1 - i\gamma_1\gamma_2), \quad \Pi_-^2 = \Pi_-, \quad [\Pi_-, (a\gamma)_\parallel] = 0.$$

The propagator was obtained in this form for the first time in Ref. [6]. Here γ_α are the Dirac matrices in the standard representation, the four-vectors with the indices $\perp$ and $\parallel$ belong to the Euclidean $\{1, 2\}$ plane and the Minkowski $\{0, 3\}$ plane correspondingly, when the field $\mathbf{B}$ is directed along the third axis. Then for arbitrary 4-vectors a_μ, b_μ one has

$$\begin{aligned} a_\perp &= (0, a_1, a_2, 0), \quad a_\parallel = (a_0, 0, 0, a_3), \\ (ab)_\perp &= (a\Lambda b) = a_1b_1 + a_2b_2, \quad (ab)_\parallel = (a\tilde{\Lambda}b) = a_0b_0 - a_3b_3, \end{aligned} \quad (4)$$

where the matrices are introduced $\Lambda_{\alpha\beta} = (\varphi\varphi)_{\alpha\beta}$, $\tilde{\Lambda}_{\alpha\beta} = (\tilde{\varphi}\tilde{\varphi})_{\alpha\beta}$, connected by the relation $\tilde{\Lambda}_{\alpha\beta} - \Lambda_{\alpha\beta} = g_{\alpha\beta} = \text{diag}(1, -1, -1, -1)$, $\varphi_{\alpha\beta} = F_{\alpha\beta}/B$ is the dimensionless tensor of the external magnetic field, $\tilde{\varphi}_{\alpha\beta} = \frac{1}{2}\varepsilon_{\alpha\beta\mu\nu}\varphi_{\mu\nu}$ is the dual tensor, the indices of four-vectors and tensors standing inside the parentheses are contracted consecutively, e.g. $(a\Lambda b) = a_\alpha\Lambda_{\alpha\beta}b_\beta$.

In spite of the translational and gauge noninvariance of the phase $\Phi(x, y)$ in the propagator (1), the total phase of three propagators in the loop is translational and gauge invariant

$$\Phi(x, y) + \Phi(y, z) + \Phi(z, x) = -\frac{e}{2}(z - x)_\mu F_{\mu\nu}(x - y)_\nu.$$

A general amplitude of the process described by Fig.1 takes the form

$$\begin{aligned} \mathcal{M} = & e^2 g_3 \int d^4 X d^4 Y Sp\{(j_3 \Gamma_3) \hat{S}(Y)(\varepsilon_2 \gamma) \hat{S}(-X-Y)(\varepsilon_1 \gamma) \hat{S}(X)\} \times \\ & \times e^{-ie(XFY)/2} e^{i(k_1 X - k_2 Y)} + (\gamma_1 \leftrightarrow \gamma_2), \end{aligned} \quad (5)$$

where $X = z - x$, $Y = x - y$, Γ_3 is the matrix corresponding to the arbitrary (S, P, V or A) vertex, g_3 is the coupling constant, j_3 is the neutrino current in the momentum space, ε_1 , k_1 and ε_2 , k_2 are the polarization vectors and the 4-momenta of initial photons.

Substituting the propagator (1), (2), and (3) into the amplitude one obtains that two parts of it which differ by the photon interchange, are proportional to the field strength B

$$\begin{aligned} \mathcal{M} \simeq & -\frac{i \alpha g_3 e B}{(4\pi)^2} \exp \left\{ -\frac{\vec{k}_{1\perp}^2 + \vec{k}_{2\perp}^2 + (\vec{k}_1 \vec{k}_2)_\perp}{2eB} \right\} \exp \left\{ -i \frac{(k_1 \varphi k_2)}{2eB} \right\} \times \\ & \times \int d^2 p Sp\{(j_3 \Gamma_3) S_\parallel(p+k_2)(\varepsilon_2 \gamma) S_\parallel(p)(\varepsilon_1 \gamma) S_\parallel(p-k_1)\} + \\ & + (\gamma_1 \leftrightarrow \gamma_2), \end{aligned} \quad (6)$$

where $S_\parallel(p) = 2\Pi_-((p\gamma)_\parallel + m)/(p_\parallel^2 - m^2)$. It should be noted, that in the amplitude (6) the projecting operators Π_- separate out the photons of only one polarization ($\perp$) of the two possible (in Adler's notation [7])

$$\varepsilon_\alpha^{(\parallel)} \sim F_{\alpha\beta} k_\beta, \quad \varepsilon_\alpha^{(\perp)} \sim \tilde{F}_{\alpha\beta} k_\beta.$$

Using the standard procedure one can transform the trace in the second term of Eq. (6) with the photon interchanged into the trace of the first term, however, with change of sign for $\Gamma_3 = P, V, A$ (and the factor $\sin[(k_1 \varphi k_2)/2eB]$ arises in the resulting amplitude) and without change of sign for $\Gamma_3 = S$ (and the factor $\cos[(k_1 \varphi k_2)/2eB]$ appears after summation). So, when the magnetic field strength is the maximal physical parameter, $eB \gg \vec{k}_\perp^2, k_\parallel^2$, only the amplitude with the scalar vertex grows linearly with the field.

The effective scalar $\nu\nu ee$ interaction arises in the standard model extensions with broken left-right (LR) symmetry [8] with mixing of the bosons

coupled with left and right charged weak currents [9]. The $\nu e W$ interaction in the model has the form

$$\begin{aligned} \mathcal{L} = & \frac{g}{2\sqrt{2}} \left\{ [\bar{e}\gamma_\alpha (1 - \gamma_5) \nu] (W_1^\alpha \cos \zeta + W_2^\alpha \sin \zeta) + \right. \\ & \left. + [\bar{e}\gamma_\alpha (1 + \gamma_5) \nu] (-W_1^\alpha \sin \zeta + W_2^\alpha \cos \zeta) + h.c. \right\}, \end{aligned} \quad (7)$$

where $W_{1,2}$ are the mass eigenstates, ζ is the mixing angle. Existing indirect limits on the LR model parameters extracted from low-energy accelerator experiments is [10]

$$M_{W_2} > 715 \text{ GeV}, \quad \zeta < 0.013. \quad (8)$$

Due to the smallness of the mixing, W_2 is almost W_R . Indirect limit on the parameters from astrophysical data (SN1987A) is even more stringent. In combination with the accelerator data it gives [11]

$$M_{W_R} > 23 \text{ TeV}, \quad \zeta < 10^{-5}. \quad (9)$$

Taking into account the smallness of the mixing angle and of the mass ratio M_{W_L}/M_{W_R} , for the effective scalar part of the $\nu\nu ee$ interaction one obtains

$$\mathcal{L}_{eff}^{(s)} \simeq -4 \zeta \frac{G_F}{\sqrt{2}} (\bar{e}e) (\bar{\nu}\nu). \quad (10)$$

With the Lagrangian (10), there exist two channels of the conversion of the photon pair into the neutrino pair, namely,

$$\gamma\gamma \rightarrow (\nu_e)_L (\bar{\nu}_e)_L, \quad \gamma\gamma \rightarrow (\nu_e)_R (\bar{\nu}_e)_R. \quad (11)$$

Here $(\nu_e)_R$ and $(\bar{\nu}_e)_L$ are the “sterile” states with respect to the standard weak interaction, which can escape from the hot and dense stellar interior.

Substitution of $\Gamma_3 = 1$, $g_3 = -4 \zeta G_F/\sqrt{2}$ and $j_3 = [\bar{\nu}(p_1)\nu(-p_2)]$, resulting from Eq. (10), into the amplitude (6) and integration give in the strong field limit

$$\begin{aligned} \mathcal{M} = & -\frac{4\alpha}{\pi} \frac{B}{B_e} \frac{\zeta G_F}{\sqrt{2} m_e} [\bar{\nu}(p_1)\nu(-p_2)] \left\{ \left(f_1^{(\perp)} f_2^{(\perp)} \right) \mathcal{F}_1 \left(\frac{q_\parallel^2}{m_e^2}, \frac{k_{1\parallel}^2}{m_e^2}, \frac{k_{2\parallel}^2}{m_e^2} \right) + \right. \\ & \left. + 4 \frac{(k_{1\parallel} f_1^{(\perp)} f_2^{(\perp)} k_{2\parallel})}{q_\parallel^2} \mathcal{F}_2 \left(\frac{q_\parallel^2}{m_e^2}, \frac{k_{1\parallel}^2}{m_e^2}, \frac{k_{2\parallel}^2}{m_e^2} \right) \right\}, \end{aligned} \quad (12)$$

here $q_{\parallel} = k_{1\parallel} + k_{2\parallel}$. Remind, that $k_{1,2\parallel} = (\omega_{1,2}, 0, 0, k_{1,2z})$. $f_{\alpha\beta}$ are the photon field tensors

$$f_{\alpha\beta}^{(\perp)} = k_{\alpha\parallel} \varepsilon_{\beta}^{(\perp)} - k_{\beta\parallel} \varepsilon_{\alpha}^{(\perp)}.$$

The functions introduced in Eq. (12) are the following

$$\begin{aligned} \mathcal{F}_1(z, s, t) &= \int_0^1 x dx \int_0^1 dy \frac{1 - 4x^2 y(1 - y)}{a^2}, \\ \mathcal{F}_2(z, s, t) &= \int_0^1 x(1 - x)(1 - 2x) dx \int_0^1 \frac{dy}{a^2}, \\ a &= 1 - zx^2 y(1 - y) - sx(1 - x)y - tx(1 - x)(1 - y). \end{aligned} \quad (13)$$

The amplitude (12) is manifestly gauge-invariant.

The cross-sections for both processes (11) are equal, $\sigma_{LL} = \sigma_{RR} \equiv \sigma$. It takes a simple form in the two limiting cases:

i) low photon energies, $\omega \lesssim m_e$

$$\sigma \simeq \frac{2\alpha^2 G_F^2 \zeta^2}{9\pi^3} \left(\frac{B}{B_e} \right)^2 \frac{k_{1\parallel}^2 k_{2\parallel}^2}{m_e^2}, \quad (14)$$

ii) high photon energies, $\omega \gg m_e$, in the leading log approximation:

$$\sigma \simeq \frac{2\alpha^2 G_F^2 \zeta^2}{\pi^3} \left(\frac{B}{B_e} \right)^2 \frac{m_e^6}{k_{1\parallel}^2 k_{2\parallel}^2} \ln^2 \frac{k_{1\parallel}^2 k_{2\parallel}^2}{m_e^4}. \quad (15)$$

The observable value in astrophysics is the stellar energy-loss from unit volume in unit time due to the neutrino escape (neutrino emissivity). In our case it can be written in the form

$$Q = \frac{1}{2} \int \frac{d^3 k_1}{(2\pi)^3} \frac{1}{e^{\omega_1/T} - 1} \int \frac{d^3 k_2}{(2\pi)^3} \frac{1}{e^{\omega_2/T} - 1} (\omega_1 + \omega_2) \frac{(k_1 k_2)}{\omega_1 \omega_2} \sigma. \quad (16)$$

Here T is the temperature of the photon gas. It is taken into account, that photons of only one polarization participate in the process, and only one (anti)neutrino from the pair can escape while the other one is trapped by the hot and dense stellar interior.

In the low temperature case, $T \lesssim m_e$, substituting (14) into (16), one obtains

$$Q_{(B)} \simeq 2.5 \cdot 10^{13} \frac{\text{erg}}{\text{cm}^3 \text{ s}} \left(\frac{\zeta}{0.013} \right)^2 \left(\frac{B}{B_e} \right)^2 \left(\frac{T}{m_e} \right)^{11}. \quad (17)$$

This should be compared with the contributions into the neutrino emissivity of the process $\gamma\gamma \rightarrow \nu\bar{\nu}$ caused by another mechanisms. For instance, the emissivity due to the finite neutrino mass is [3]

$$Q_{(m_\nu)} \simeq 0.4 \cdot 10^5 \frac{\text{erg}}{\text{cm}^3 \text{ s}} \left(\frac{m_\nu}{1 \text{ eV}} \right)^2 \left(\frac{T}{m_e} \right)^{11}. \quad (18)$$

On the other hand, in the case of non-locality of the weak interaction, investigated in Ref. [4], one can estimate the emissivity, which is suppressed by the factor $(m_e/M_W)^4$:

$$Q_{(NL)} \sim 10 \frac{\text{erg}}{\text{cm}^3 \text{ s}} \left(\frac{T}{m_e} \right)^{13}. \quad (19)$$

It is seen that for $B \gtrsim B_e$, and for mixing at the level of $\zeta \sim 10^{-5}$, the field-induced mechanism of the reaction $\gamma\gamma \rightarrow \nu\bar{\nu}$ strongly dominates all the other mechanisms.

In the high temperature case, $T \gg m_e$, substituting (15) into (16), one obtains

$$Q_{(B)} \simeq 0.4 \cdot 10^{12} \frac{\text{erg}}{\text{cm}^3 \text{ s}} \left(\frac{\zeta}{0.013} \right)^2 \left(\frac{B}{B_e} \right)^2 \left(\frac{T}{m_e} \right)^3 \ln^5 \frac{T}{m_e}. \quad (20)$$

In order to make a numerical estimation, let us consider the Supernova explosion with generation of very strong magnetic field $B \sim 10^3 B_e$, see e.g. [12], with the temperature $T \sim 35 \text{ MeV}$ which is believed to be typical for the Supernova core [13], and $V \sim 10^{18} \text{ cm}^3$. For the contribution of the considered field-enhanced process $\gamma\gamma \rightarrow \nu\bar{\nu}$ into the neutrino luminosity we obtain

$$\frac{dE}{dt} \sim 2 \cdot 10^{44} \frac{\text{erg}}{\text{sec}} \left(\frac{\zeta}{0.013} \right)^2. \quad (21)$$

It is too small if compared with the typical Supernova neutrino luminosity 10^{52} erg/sec.

In summary, we have performed a general analysis of the three-vertex loop in a strong magnetic field and have considered the photon-neutrino process $\gamma\gamma \rightarrow \nu\bar{\nu}$. It is shown that various types of the neutrino - electron effective interactions lead to different dependences of the amplitudes on the field strength. The effective scalar $\nu\bar{\nu}ee$ coupling which exists in the extensions of the standard model with broken left-right symmetry, leads to the enhancement of the process $\gamma\gamma \rightarrow \nu\bar{\nu}$ by strong external magnetic field. For the field strength $B \gtrsim B_e$ this mechanism could dominate other mechanisms of the process $\gamma\gamma \rightarrow \nu\bar{\nu}$ in the neutrino emissivity. However, its contribution into the Supernova energetics is rather small.

This work was supported in part by the Russian Foundation for Basic Research under the Grant N 01-02-17334 and by the Ministry of Education of Russian Federation under the Grant No. E00-11.0-5.

References

- [1] M. Gell-Mann, *Phys. Rev. Lett.* **6** (1961) 70.
- [2] R. J. Crewther, J. Finjord and P. Minkowski, *Nucl. Phys.* **B 207** (1982) 269.
- [3] S. Dodelson and G. Feinberg, *Phys. Rev.* **D 43** (1991) 913.
- [4] D. A. Dicus and W. W. Repko, *Phys. Rev.* **D 48** (1993) 5106.
- [5] R. Shaisultanov, *Phys. Rev. Lett.* **80** (1998) 1586; T.-K. Chyi et al., *Phys. Lett.* **B 466** (1999) 274; D. A. Dicus and W. W. Repko, *Phys. Lett.* **B 482** (2000) 141.
- [6] V. V. Skobelev, *Izv. Vuz., Fiz.* **10** (1975) 142; Yu. M. Loskutov and V. V. Skobelev, *Phys. Lett.* **A 56** (1976) 151.
- [7] S. L. Adler, *Ann. Phys. N.Y.* **67** (1971) 599.
- [8] E. M. Lipmanov, *Yader. Fiz.* **6** (1967) 571 [*Sov. J. Nucl. Phys.* **6** (1968) 395]; J. C. Pati and A. Salam, *Phys. Rev.* **D 10** (1974) 275.

- [9] M. A. B. Bég, R. Budny, R. N. Mohapatra and A. Sirlin, *Phys. Rev. Lett.* **38** (1977) 1252.
- [10] D. E. Groom e. a. (Particle Data Group), *Eur. Phys. J. C* **15** (2000) 1.
- [11] R. Barbieri and R. N. Mohapatra, *Phys. Rev. D* **39** (1989) 1229.
- [12] G. S. Bisnovatyi-Kogan, *Astro. Zh.* **47** (1970) 813 [*Sov. Astron.* **14** (1971) 652]; R. C. Duncan and C. Thompson, *Astrophys. J.* **392** (1992) L9.
- [13] G. G. Raffelt, *Stars as Laboratories for Fundamental Physics*, University of Chicago Press (Chicago, 1996).




Fig. 1